

Bibliografia

Aldo Franco de Rose; Fabrizio Gallo; Matheo Giglio and Georgio Carnignani. Role of mepartricine in category III chronic non bacterial prostatitis/chronic pelvic pain sindrome: a randomized prospective placebo-controlled trial. Urology 63:13-19, 2004

British National Formulary (BNF), September 2013 – March 2014

Cunningham Gr, Settel AD.Treament of male sexual dysfunction. Uptodate 2014

European Association of Urology - Guidelines 2014

EAU guidelines on urinary incontinence 2014; EAU guidelines on the management of non-neurogenic male lower urinary tract symptoms (LUTS), incl benign prostatic obstruction 2014; EAU guidelines on neuro-urology 2014; Diagnosis and treatment of overactive bladder (non-neurogenic) in adults - AUA guideline 2014; The effects of anti-muscarinic treatments in overactive bladder: na update of a systematic review and meta-analysis Chapple et al. Eur Urol 54 (2008), 543-562; The role of antimuscarinics in the managment of men with symptoms of overactive bladder associated with concomitant bladder outlet obstructon an update Athanasopoulos et al. Eur Urol 60(2011), 94-105

EAU guidelines on urinary incontinence 2014; Diagnosis and treatment of overactive bladder (non-neurogenic) in adults - AUA guideline 2014; Comparative Efficacy and Safety of Medical Treatments for the Management of Overactive Bladder: A Systematic Literature Review and Mixed Treatment Comparison. Eur Urol 65 (2014), 755-765; A randomized controlled study to evaluate the efficacy of tamsulosin monotherapy and its combination with mirabegron on patients with overactive bladder induced by benign prostatic obstruction. J Urol. 2014 Sep 21. pii: S0022-5347(14)04488-7. doi: 10.1016/j.juro.2014.09.091. [Epub ahead of print]; Urodynamics and safety of the β_3 -adrenoceptor agonist mirabegron in males with lower urinary tract symptoms and bladder outlet obstruction. J Urol. 2013 Oct;190(4):1320-7

EAU guidelines on urinary incontinence 2014; EAU guidelines on the management of non-neurogenic male lower urinary tract symptoms (LUTS), incl benign prostatic obstruction 2014; EAU guidelines on neuro-urology 2014; EAU guidelines on cronic pelvic pain 2014; Diagnosis and treatment of overactive bladder (non-neurogenic) in adults - AUA guideline 2014; An Updated Systematic Review and Statistical Comparison of Standardised Mean Outcomes for the Use of Botulinum Toxin in the Management of Lower Urinary Tract Disorders. Eur Urol 65 (2014), 981-990

Kunelius P, Hakkinen J, Wkkarinen O.Is high-dose Yohimbine hydrochloride effective in the treatment of mixed-type impotence? A prospective, randomized, controlled double-blind crossover study. Urology 49:441-444.1997